

# Sodium feredetate for the treatment of anaemia

A separate leaflet about the use of sodium feredetate for the prevention of anaemia is available [www.medicinesforchildren.org.uk](http://www.medicinesforchildren.org.uk).



This leaflet is for parents and carers about how to use this medicine in children. Our information sometimes differs from that provided by the manufacturers, because their information is usually aimed at adults. Please read this leaflet carefully. Keep it somewhere safe so that you can read it again.

## Name of medicine

### Sodium feredetate

**Brand name:** Sytron, SodiFer

## Why is it important for my child to take this medicine?

If the body has too little iron (iron deficiency), it cannot make enough haemoglobin, which it needs to carry oxygen in the blood around the body. Having too little haemoglobin is called anaemia. Children with anaemia are often pale, feel tired and have little energy, and may not grow or develop properly. sodium feredetate contains lots of iron. This helps the body to make more haemoglobin and so treats the anaemia.

## What is sodium feredetate available as?

- **Liquid medicine:** 190 mg (which is 27.5 mg iron) in 5 mL; this medicine contains a small amount of alcohol


## When should I give sodium feredetate?

- Sodium feredetate is usually given **three times** a day. This should be first thing in the morning, early afternoon and at bedtime. Ideally, these times are at least 6 hours apart, for example 8 am, 2 pm and 8 pm.
- In some circumstances sodium feredetate may be given just **one or two times** a day. If this applies to your child then your doctor will explain this to you.

Give the medicine at about the same time each day so that this becomes part of your child's daily routine, which will help you to remember.

## How much should I give?

Your doctor will work out the amount of sodium feredetate that is right for *your* child. The dose will be shown on the medicine label.

 **It is important that you follow your doctor's instructions about how much to give.**

## How should I give it?

Do not give the medicine with a milk feed or with food containing milk or dairy products (e.g. cheese or yogurt).



**Liquid medicine:** Measure out the right amount an oral syringe or medicine spoon. You can get these from your pharmacist. Do not use a kitchen teaspoon as it will not give the right amount.

## When should the medicine start working?

Sodium feredetate takes some time to work. You might notice an improvement in your child when they have been taking the medicine for 3–4 weeks – they may be less pale, have more energy and have less shortness of breath. Your child will need to take sodium feredetate regularly for at least 3 months for it to have its full benefits.

## What if my child is sick (vomits)?

- If your child is sick less than 30 minutes after having a dose of sodium feredetate, give them the same dose again.
- If your child is sick more than 30 minutes after having a dose of sodium feredetate, you **do not** need to give them another dose. Wait until the next normal dose.

If your child is sick again, seek advice from your GP, pharmacist or hospital.

## What if I forget to give it?

If you miss a dose, wait until the next normal dose. Do not give the missed dose.



**Never give a double dose of sodium feredetate.**

## What if I give too much?



It can be dangerous to give too much sodium feredetate. Never give your child more than the doctor has advised.

If your child has one or more of the following symptoms, they may have had too much sodium feredetate:

- stomach pains
- being repeatedly sick (vomiting)
- diarrhoea
- the vomit or stools (poo) may be blood stained, or be green or grey.



Children who have had too much sodium feredetate might not show any symptoms, or only have mild symptoms. If you think you may have given your child too much, contact your doctor or local NHS services (details at end of leaflet) or take your child to hospital.



If your child appears very unwell or drowsy call for an ambulance **straight away**.

Take the medicine container or packaging with you, even if it is empty. This will be useful to the doctor. Have the medicine or packaging with you if you telephone for advice.

## Are there any possible side-effects?

We use medicines to make our children better, but sometimes they have other effects that we don't want (side-effects).

- When your child first starts having Sytron® they may have stomach pain, feel sick or be sick (vomit) or they

may get diarrhoea, wind (flatulence) or constipation (difficulty doing a poo). It may help to give each dose after some food even though iron preparations, like sodium feredetate, are best absorbed on an empty stomach. If these problems last longer than a week, contact your doctor.

- Your child's stool (poo) may look darker than usual. This is nothing to worry about.

There may, sometimes, be other side-effects that are not listed above. If you notice anything unusual and are concerned, contact your doctor. You can report any suspected side-effects to a UK safety scheme at <http://www.mhra.gov.uk/yellowcard>.

### Can other medicines be given at the same time as sodium feredetate?

- You can give your child medicines that contain paracetamol or ibuprofen, unless your doctor has told you not to.
- Sodium feredetate should not be given with any other medicines or vitamin preparations that contain iron.
- Try not to give the sodium feredetate at the same time as a milk feed or dairy products (e.g. cheese or yogurt), as less of the medicine will be absorbed into the body.
- Sodium feredetate should not be taken with some medicines that you get on prescription. Tell your doctor and pharmacist about any other medicines your child is taking before giving sodium feredetate.
- Check with your doctor or pharmacist **before** giving any other medicines to your child. This includes herbal or complementary medicines.

### Is there anything else I need to know about this medicine?

- A separate leaflet about the use of sodium feredetate for the treatment of anaemia is available on the Medicines for Children website, [www.medicinesforchildren.org.uk](http://www.medicinesforchildren.org.uk).
- Once opened, this medicine should not be kept for longer than 3 months.

### General advice about medicines

- Try to give medicines at about the same times each day, to help you remember.
- If you are not sure a medicine is working, contact your doctor but continue to give the medicine as usual in the meantime. Do not give extra doses, as you may do harm.

- Only give this medicine to *your* child. Never give it to anyone else, even if their condition appears to be the same, as this could do harm.



If you think someone else may have taken the medicine by accident, contact your doctor straight away.

- Make sure that you always have enough medicine. Order a new prescription at least 2 weeks before you will run out.
- Make sure that the medicine you have at home has not reached the 'use by' date on the packaging. Give old medicines to your pharmacist to dispose of.

### Where should I keep this medicine?

- Keep the medicine in a cupboard, away from heat and direct sunlight. It does not need to be kept in the fridge.
- Make sure that children cannot see or reach the medicine.
- Keep the medicine in the container it came in.

### Who to contact for more information

Your doctor, pharmacist or nurse will be able to give you more information about sodium feredetate and other medicines used to prevent or treat anaemia.

You can also get useful information from:

**England: NHS 111**

Tel 111 - [www.nhs.uk](http://www.nhs.uk)

**Scotland: NHS 24**

Tel 111 - [www.nhs24.scot](http://www.nhs24.scot)

**Wales: NHS 111 Wales**

Tel 111 - [www.111.wales.nhs.uk](http://www.111.wales.nhs.uk)

**Northern Ireland: NI Direct**

[www.nidirect.gov.uk](http://www.nidirect.gov.uk)

[www.medicinesforchildren.org.uk](http://www.medicinesforchildren.org.uk)